SN. 09/965,796

ATTORNEY DOCKET No. IMMU:007US3

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

- 24. (Currently amended) A method for treating a subject having a B-cell malignancy, comprising administering to the subject a therapeutic composition comprising a pharmaceutically acceptable carrier, and an immunoconjugate, wherein the immunoconjugate comprises
- (i) at least one human, humanized or chimeric anti-CD22 antibody or a fragment thereof, wherein the antibody fragment is selected from the group consisting of F(ab')2, Fab', Fab, and single chain Fv, and
- (ii) a drug or a radioisotope, wherein said radioisotope is other than ¹³¹I, wherein the immunoconjugate is used in combination with a naked anti-CD2O mAb.
- 25. (Previously presented) The method according to claim 24, wherein the immunoconjugate comprises a chemotherapeutic drug.
- 26. (Previously presented) The method according to claim 25, wherein the chemotherapeutic drug is selected from the group consisting of cyclophospharnide, etoposide, vincristine, procarbazine, prednisone, carmustine, doxorubicin, methotrexate, bleomycin, dexamethasone, phenyl butyrate, bryostatin-1 and leucovorin.
- 27. (Previously presented) The method according to claim 25, wherein the chemotherapeutic drug is selected from the group consisting of nitrogen mustards, alkyl sulfonates, nitrosoureas, triazenes, folic acid analogs, pyrimidine analogs, purine analogs, antibiotics, epipodophyllotoxins, platinum coordination complexes, and hormones.

28-35. (Canceled)

- 36. (Currently amended) The method according to claim 24, wherein the anti-CD22 antibody or fragment thereof is a human antibody or antibody fragment.
- 37. (Currently amended) The method according to claim 24, wherein the anti-CD22 antibody or fragment thereof is a humanized antibody or antibody fragment.

SN. 09/965,796

ATTORNEY DOCKET No. IMMU:007US3

- 38. (Currently amended) The method according to claim 24, wherein the anti-CD22 antibody er fragment thereof is a chimeric antibody or antibody fragment.
- 39. (Currently amended) The method according to claim 24, wherein the anti-CD22 antibody or fragment thereof comprises a multivalent fusion protein that additionally comprises at least one antibody component that binds with CD19, CD2O, CD52 or CD74.
- 40. (Currently amended) The method according to claim 39, wherein the anti-CD22 antibody or fragment thereof comprises a trivalent fusion protein.
- 41, (Currently amended) The method according to claim 39, wherein the anti-CD22 antibody or fragment thereof comprises a tetravalent fusion protein.
- 42. (Currently amended) The method according to claim 39, wherein the anti-CD22 antibody or fragment thereof comprises a quintavalent fusion protein.
- 43. (Currently amended) The method according to claim 24, wherein the immunoconjugate comprises polyethyleneglycol to extend the half-life of the antibody or fragment thereof, in blood, lymph, or other extracellular fluids.
- 44. (Currently amended) The method according to claim 24, wherein the anti-CD22 antibody or antibody fragment is a human antibody or antibody fragment.
 - 45. (Canceled)
 - 46. (Canceled)
- 47. (Previously presented) The method according to claim 24, wherein the therapeutic composition comprises at least two monoclonal antibodies that bind with distinct CD22 epitopes, wherein the CD22 epitopes are selected from the group consisting of epitope A, epitope B, epitope C, epitope D and epitope E.

48-51. (Canceled)

52. (Previously presented) The method according to claim 24, wherein the radioisotope is selected from the group consisting of ⁹⁸Au, ³²P, ¹²⁵I, ⁹⁰Y, ¹⁸⁶Re, ⁸⁸Re, ⁶⁷Cu, ²¹¹At, ²¹³Bi, and ²²⁴Ac

: .

SN. 09/965,796

ATTORNEY DOCKET No. IMMU:007US3

53-54. (Canceled)

- (Previously presented) The method according to claim 24, wherein the anti-CD22 **55**. immunoconjugate comprises a ⁹⁰Y radioisotope.
- (Previously presented) The method according to claim 55, wherein the 90Y is 56. attached to the anti-CD22 immunoconjugate by means of chelating agent.
- (Previously presented) The method according to claim 56, wherein the chelating 57. agent is diethylenetriaminepentaacetic acid.
- (Previously presented) The method according to claim 38, wherein the radioisotope 58. is ⁶⁷Cu.
- (Previously presented) The method according to claim 58, wherein the chelating 59. agent is p-bromoacetamido-benzyl-tetraethylaminetetraacetic acid.

60-97. (Canceled)